This Page Is Inserted by IFW Operations and is not a part of the Official Record

BEST AVAILABLE IMAGES


Defective images within this document are accurate representations of the original documents submitted by the applicant.


Defects in the images may include (but are not limited to):


- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.


As rescanning documents will not correct images, please do not report the images to the Image Problem Mailbox.







Cell attachment Heparanase T T 1E 1E 2E 2E kDa - + - + - + Cb Cb Cc



T = Trypsin

1E = 1mM EDTA


2E = 2mM EDTA

Cb = Control, purified heparanase from baculovirus, p60

Cc = Control, purified heparanase from CHO cells, p45

Fig. 2b

Heparanase adherence to cells requires the presence of GAGs



Sample O.D. - sample O.D. - blank Δ O.D. B16-F1 0.1245 0.0422 0.0823 B16-F1 transfected 0.1671 0.0303 0.1366 B16-F1 adhered 0.4448 0.0321 0.4127 Control b22 0.2258 0.1163 0.1095

Fig. 4a



b22, b27, bmix = control, purified baculo heparanase (p60), different batches.
c = control, purified CHO heparanase (p45)
C = Control, substrate only

Fig. 4b




Fig. 4c

Sample #	Time of incubation with heparanase	Cell detachment from culture plate	O.D sample	Δ O.D.
1	0	Trypsin	0.1291	
2	15 min	Trypsin	0.1777	0.0486
3	60 min	Trypsin	0.318	0.1847
4	180 min	Trypsin	0.3633	0.2342
5	0	1mM EDTA	0.1370	
6	180 min	1mM EDTA	0.3754	0.2384

Fig. 5a




Fig. 5b




Fig. 6a





Fig. 6b



Cb = Control, from baculovirus, p60. **Cc** = Control, from CHO cells, p45.

Fig. 6c






Fig. 7








Fig. 8e

